

Original Article

Structure-based design of functionalized 2-substituted and 1,2-disubstituted benzimidazole derivatives and their *in vitro* antibacterial efficacy


Olayinka O. Ajani^{a,*}, Olayinka O. Tolu-Bolaji^a, Shade J. Olorunshola^b, Yuxia Zhao^c,
Damilola V. Aderohunmu^a

^a Department of Chemistry, C.S.T., Covenant University, Canaanland, km 10, Idiroko Road, P.M.B. 1023, Ota, Ogun State, Nigeria

^b Department of Biological Sciences, C.S.T., Covenant University, Canaanland, km 10, Idiroko Road, P.M.B. 1023, Ota, Ogun State, Nigeria

^c Technical Institute of Physics and Chemistry, CAS, No 29, Zhongguancun East Road, Haidian District, Beijing 100190, China

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 23 June 2017

Revised 15 September 2017

Accepted 20 September 2017

Available online 22 September 2017

Keywords:

Benzimidazole

Antibacterial

Activity index

Cycloaddition

Spectroscopy

ABSTRACT

The aim of this present study was to synthesize 2-substituted and 1,2-disubstituted benzimidazole derivatives to investigate their **antibacterial** diversity for possible future drug design. The structure-based design of precursors 2-(1H-benzimidazol-2-yl)aniline **1**, 2-(3,5-dinitro phenyl)-1H-benzimidazole **3** and 2-benzyl-1H-benzimidazole **5** were achieved by the condensation reaction of *o*-phenylenediamine with anthranilic acid, 3,5-dinitrophenylbenzoic acid, and phenylacetic acid, respectively. The precursors **1**, **3** and **5**, upon reaction with six different electrophile-releasing agents, furnished the corresponding 2-substituted benzimidazole, **2a-f** and 1,2-disubstituted benzimidazole derivatives **4a-f** and **6a-f**, respectively. The structural identity of the targeted compounds was authenticated by elemental analytical data and spectral information from FT-IR, UV, ¹H, and ¹³C NMR. The outcome of the findings from the *in vitro* screening unveiled 2-benzyl-1-(phenylsulfonyl)-1H-benzimidazole **6b** as the most active derivative with lowest MIC value of 15.63 µg/mL.

© 2017 Production and hosting by Elsevier B.V. on behalf of Cairo University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

From time to time, heterocyclic templates have continued to gain respect and much interest among the medicinal chemists, because of their numerous therapeutic applications and effective reported druggability [1]. Benzimidazole is a heterocyclic aromatic

Peer review under responsibility of Cairo University.

* Corresponding author.

E-mail address: ola.ajani@covenantuniversity.edu.ng (O.O. Ajani).

<https://doi.org/10.1016/j.jare.2017.09.003>

2090-1232/© 2017 Production and hosting by Elsevier B.V. on behalf of Cairo University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

organic compound that plays important functions in the development of theory in heterocyclic chemistry and organic synthesis [2]. Benzimidazole is a strongly acidic compound with a pKa of 12.75, while its conjugated acid has a pKa of 5.68, which is less basic than imidazole. Benzimidazole is readily prepared by [4+1]-cycloaddition of *o*-phenylenediamine with a one carbon donor source in the presence of various heterogeneous catalysts [3], such as H₂O₂/HCl [4], H₂O₂/CAN [5], H₂O/HCl [6], H₂O₂/Fe(NO₃)₃ [7], and H₂O₂/Bu₄NI [8] as efficient oxidative couples. In healthful analysis, the synthesis of novel benzimidazole derivatives remains a focus [9]. Diverse synthetic efforts for accessing benzimidazole derivatives have been documented, however, the commonest technique involves the reaction of *o*-phenylenediamine with alkanic acids. From the evaluation of the works of various researchers, benzimidazole derivatives have been reported to possess antimalarial [10], anticancer [11], antimicrobial [12,13], antioxidant [14], and anticonvulsant [15] activities among others. Some derivatives of benzimidazole are well known in corrosion studies and their corrosion inhibition efficiencies are related to their adsorption properties [16].

The outbreak of new diseases and the increase in population of drug resistant strains of bacteria, such as methicillin-resistant *Staphylococcus aureus* [17], vancomycin-resistant *Enterococci* [18], ampicillin-resistant *Enterobacter aerogenes* [19], gentamicin-resistant *Escherichia coli* [20], and chloroquine-resistant *Plasmodium falciparum* [21], have posed great challenges to life and wellbeing of mankind. Based on the existence of antidrug multi-resistant bacteria strains [22], the occurrence of side effects to commercially available drugs [23], adverse drug reaction in elderly patient [24], the emergence of new diseases, and global health threat that have resulted in high mortality rate [25]; it has become highly imperative to consistently and continuously engage in the synthetic preparation of novel heterocyclic templates as highly dynamic biologically active substances for therapeutic uses. Therefore, it is beneficial to design some 2-substituted- and 1,2-disubstituted benzimidazole derivatives by ecofriendly method so as to examine their antimicrobial properties for possible future drug development.

Material and methods

Chemical compounds and reagents were purchased from Sigma-Aldrich Chemicals (St. Louis, Missouri, USA) apart from Tetrahydrofuran (THF), benzenesulfonyl chloride, and anthranilic acid which were supplied by the British Drug Houses (Poole, Dorset, England). All these compounds were then made available by Department of Chemistry, Covenant University for research use. All the chemicals are pure and they were used directly without further purification. The synthesized heterocyclic frameworks were evaluated for their melting point determination using Stuart equipment and the value obtained were recorded directly. Bruker fourier-transform (ft-ir) spectrophotometer was utilized to obtain infrared data. The UV spectra of the solution of the compounds in THF were run in UV Genesys 10 s. The ¹H and ¹³C nuclear magnetic resonance of the heterocycles were NMR Bruker DPX 400 spectrometer at 400 MHz and 100 MHz, respectively in DMSO-*d*₆. The reference utilized was Tetramethylsilane (TMS). The reaction progress as well as the level of purity was routinely checked and monitored with Thin Layer Chromatography (TLC) using CHCl₃/CH₃OH (9:1, v/v) eluent. After reaction was completed, solvents were evaporated under reduced pressure using IKA® RV 10 Rotary evaporator. In a situation where more than one spots were observed, column chromatography was carried out to get a pure compound.

2-(1H-Benzimidazol-2-yl)aniline as precursor 1

o-Phenylene diamine (15.00 g, 140.00 mmol) was weighed and dissolved in 150 mL of ethanol in a round-bottomed flask. It was stirred for 5 min with the aid of magnetic stirrer after which anthranilic acid (19.20 g, 140.00 mmol) was gradually tipped into the solution followed by the addition of a catalytic amount of NH₄-Cl (0.75 g, 14.00 mmol). The resulting solution was then heated under reflux at 60–70 °C for 2 h. The TLC was utilized to ascertain the progress of reaction. Upon completion, the resulting solution was allowed to cool down. The flask content was evaporated to dryness and triturated with ice-cold water. The solid mass formed was separated by suction filtration to furnish 2-(1H-benzimidazol-2-yl)aniline **1** (72.68%), mp = 85–87 °C, colour = gray. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 4.50 (s-br, 2H, NH₂), 6.37–6.39 (dd, *J*₁ = 4.54 Hz, *J*₂ = 8.80 Hz, 1H, Ph-H), 6.49–6.50 (dd, *J*₁ = 4.21 Hz, *J*₂ = 8.87 Hz, 1H, Ph-H), 6.84–6.86 (d, *J* = 8.80 Hz, 1H, Ph-H), 7.12–7.14 (d, *J* = 8.00 Hz, 1H, Ph-H), 7.39–7.41 (m, 2H, Ph-H), 7.59–7.63 (d, *J* = 8.87 Hz, 1H, Ph-H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 109.6, 115.2 (2 × CH), 116.9, 119.4, 123.1 (2 × CH), 125.5, 129.7, 141.9 (2 × C), 145.1, 155.0 ppm. λ_{max} in nm (log ε_{max}): 218 (4.2741), 253 (4.3096), 326 (3.6434). FT-IR ν in cm⁻¹: 3424 (N–H of NH₂), 3422 (N–H of NH₂), 3405 (N–H), 1620 (C=C aromatic). Anal. Calcd for C₁₃H₁₁N₃ (209.25): C, 74.62; H, 5.30; N, 20.08%. Found: C, 74.80; H, 5.47; N, 19.96%.

Overall protocol towards accessing 2-substituted benzimidazole 2a-f

Precursor **1** (4.00 g, 19.10 mmol) was dissolved in 20 mL of tetrahydrofuran (THF) in a round-bottomed flask at room temperature. The medium was basified by the addition of Na₂CO₃ (4.06 g, 38.30 mmol) and cooled to 0–5 °C in ice bath. The corresponding electrophile-releasing substrate **a-f** (19.10 mmol) was then added and the reacting mixture was maintained on ice bath for additional 15 min after which the medium was warmed up to room temperature and stirred there for 24 h. Monitoring of reaction progress was conducted using TLC and upon reaction completion, the solvent was evaporated at reduced pressure using rotary evaporator. Cold water was added to the resulting mass, filtered by suction, and air-dried to afford crude product which upon column purification afforded 2-substituted benzimidazole derivatives **2a-f**.

N-(2-(1H-Benzimidazole-2-yl)phenyl) acetamide 2a

When **a** = acetyl chloride, yield 57.70%, mp = 253–255 °C, colour = gray. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 2.67 (s, 3H, CH₃–CO), 6.36–6.39 (dd, *J*₁ = 4.58 Hz, *J*₂ = 8.82 Hz, 1H, Ph-H), 6.51–6.52 (dd, *J*₁ = 4.18 Hz, *J*₂ = 8.89 Hz, 1H, Ph-H), 6.86–6.88 (d, *J* = 8.82 Hz, 1H, Ph-H), 7.13–7.15 (d, *J* = 8.00 Hz, 1H, Ph-H), 7.39–7.43 (m, 2H, Ph-H), 7.62–7.65 (d, *J* = 8.89 Hz, 1H, Ph-H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 26.4 (CH₃), 109.6, 115.2 (2 × CH), 116.8, 119.6, 123.6 (2 × CH), 125.6, 129.8, 141.7 (2 × C), 145.3, 156.1, 175.4 (C=O) ppm. λ_{max} in nm (log ε_{max}): 218 (4.2988), 248 (4.5563), 323 (3.8261), 464 (2.4771). FT-IR ν in cm⁻¹: 3405 (N–H), 1699 (C=O). Anal. Calcd for C₁₅H₁₃N₃O (251.11): C, 71.70; H, 5.21; N, 16.72%. Found: C, 71.88; H, 5.09; N, 16.89%.

N-(2-(1H-Benzimidazole-2-yl)phenyl)benzenesulfonamide 2b

When **b** = benzenesulfonyl chloride, yield 97.21%, mp = N.D. (Oily), colour = black. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 6.35–6.38 (m, 3H, Ph-H), 6.46–6.49 (m, 2H, Ph-H), 6.84–6.86 (d, *J* = 7.16 Hz, 2H, Ph-H), 7.11–7.14 (d, *J* = 11.96 Hz, 2H, Ph-H), 7.40–7.42 (m, 2H, Ph-H), 7.62–7.65 (d, *J* = 11.88 Hz, 2H, Ph-H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 109.8, 115.3 (2 × CH), 116.9, 119.5, 123.2 (2 × CH), 125.6, 127.5 (2 × CH), 128.9, 129.8

(2 × CH), 131.9, 154.8, 139.7, 141.6 (2 × C), 145.3 ppm. λ_{max} in nm (log ϵ_{max}): 221 (4.6609), 251 (4.7332), 317 (4.2878), 428 (3.9294). FT-IR ν in cm^{-1} : 3405 (N–H), 3266 (N–H), 1620 (C=C aromatic), 1575 (C=N imine), 1376 (SO_2), 1185 (SO_2). Anal. Calcd for $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_2\text{S}$ (349.09): C, 65.31; H, 4.33; N, 12.03%. Found: C, 65.20; H, 4.15; N, 11.83%.

N-(2-(1*H*-Benzimidazole-2-yl)phenyl)-4-methylbenzenesulfonamide **2c**

When **c** = *p*-toluenesulfonyl chloride, yield 80.03%, mp = N.D. (Oily), colour = brown. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 2.80 (s, 3H, CH_3 -Ar), 6.36–6.39 (m, 3H, Ph–H), 6.49–6.51 (m, 2H, Ph–H), 6.84–6.86 (d, J = 7.04 Hz, 1H, Ph–H), 7.11–7.14 (d, J = 11.96 Hz, 2H, Ph–H), 7.40–7.42 (m, 2H, Ph–H), 7.62–7.65 (d, J = 11.88 Hz, 2H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 21.3 (CH_3), 109.8, 115.3 (2 × CH), 116.9, 119.5, 123.2 (2 × CH), 125.6, 127.3 (2 × CH), 129.1, 129.8 (2 × CH), 131.9, 139.7, 141.8 (2 × C), 145.3, 155.0 ppm. λ_{max} in nm (log ϵ_{max}): 212 (4.6343), 233 (5.2124), 311 (4.6750). FT-IR ν in cm^{-1} : 3407 (N–H), 3263 (N–H), 1377 (SO_2), 1187 (SO_2). Anal. Calcd for $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$ (363.43): C, 66.10; H, 4.71; N, 11.56%. Found: C, 65.95; H, 4.63; N, 11.75%.

2-(1*H*-Benzimidazol-2-yl)-*N*-(3-chlorobenzyl)aniline **2d**

When **d** = 3-chlorobenzyl chloride, yield 87.90%, mp = N.D. (Oily), colour = black. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 3.73 (s, 2H, CH_2), 6.31–6.35 (m, 3H, Ph–H), 6.48–6.50 (m, 2H, Ph–H), 6.84–6.87 (d, J = 8.00 Hz, 1H, Ph–H), 7.11–7.14 (d, J = 11.96 Hz, 2H, Ph–H), 7.61–7.63 (d, J = 7.96 Hz, 2H, Ph–H), 7.92–7.94 (d, J = 7.72 Hz, 1H, Ph–H), 8.21 (s, 1H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 48.2 (CH_2), 112.3, 114.1, 115.1 (2 × CH), 117.4, 123.2 (2 × CH), 125.0, 125.4, 126.1, 126.8, 129.3, 129.9, 132.2, 134.3, 141.8 (2 × C), 145.4, 154.8 ppm. λ_{max} in nm (log ϵ_{max}): 224 (5.0026), 251 (4.9633), 302 (4.7279). FT-IR ν in cm^{-1} : 3460, 3354 (N–H), 3107 (C–H aromatic), 2924 (CH aliphatic), 2854 (CH aliphatic), 1624 (C=C Aromatic), 1581 (C=N). Anal. Calcd for $\text{C}_{20}\text{H}_{16}\text{N}_3\text{Cl}$ (333.81): C, 71.96; H, 4.84; N, 12.59%. Found: C, 72.14; H, 5.02; N, 12.38%.

5-((2-(1*H*-Benzimidazole-2-yl)phenyl)amino)pyrimidine-2,4(1*H*,3*H*)-dione **2e**

When **e** = 5-bromouracil, yield 83.29%, mp > 300 °C, colour = - brown. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 6.36–6.39 (m, 1H, Ph–H), 6.50–6.52 (m, 1H, Ph–H), 6.83–6.86 (d, J = 11.88 Hz, 1H, Ph–H), 7.11–7.13 (d, J = 8.00 Hz, 2H, Ph–H), 7.39–7.43 (m, 2H, Ph–H), 7.62–7.65 (d, J = 11.88 Hz, 1H, Ph–H), 8.37–8.41 (d, J = 13.92 Hz, 1H, Ph–H), 11.02 (s, 1H, NH), 11.56 (s, 1H, NH). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 109.2, 110.6, 115.6 (2 × CH), 119.3, 123.1 (2 × CH), 125.3, 125.7, 126.2, 129.4, 141.8 (2 × C), 145.0, 152.5, 169.0 (C=O), 169.8 (C=O) ppm. λ_{max} in nm (log ϵ_{max}): 202 (4.5646), 224 (4.9854), 254 (5.1392). FT-IR (ν_{max} in cm^{-1}): 3460, 3354 (N–H), 3107 (C–H aromatic), 2924 (CH aliphatic), 2854 (CH aliphatic), 1624 (C=C aromatic), 1581 (C=N). Anal. Calcd for $\text{C}_{17}\text{H}_{13}\text{N}_5\text{O}_2$ (319.32): C, 63.94; H, 4.10; N, 21.93%. Found: C, 63.90; H, 3.99; N, 22.01%.

N-(2-(1*H*-Benzimidazole-2-yl)phenyl)benzene-1,4-diamine **2f**

When **f** = 4-chloroaniline, yield 80.74%, mp > 300 °C, colour = - gray. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 5.32 (s, 2H, NH_2), 6.33–6.38 (m, 2H, Ph–H), 6.49–6.53 (m, 2H, Ph–H), 6.83–6.86 (d, J = 9.92 Hz, 1H, Ph–H), 7.11–7.14 (d, J = 10.48 Hz, 2H, Ph–H), 7.39–7.41 (d, J = 8.50 Hz, 2H, Ph–H), 7.59–7.62 (d, J = 10.48 Hz, 2H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 109.8, 115.3

(2 × C), 117.2 (2 × CH), 118.1, 119.5, 121.1 (2 × CH), 123.0 (2 × CH), 125.6, 129.8, 132.2, 137.9, 140.9, 141.8 (2 × C), 154.8 ppm. λ_{max} in nm (log ϵ_{max}): 224 (5.0228), 254 (1.9881), 308 (4.5453). FT-IR (ν_{max} in cm^{-1}): 3436 (N–H of 1° amine), 3406 (N–H of 1° amine), 3330 (N–H of 2° amine), 3060 (C–H aromatic), 1613 (C=C aromatic), 1577 (C=N). Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{N}_4$ (300.36): C, 75.98; H, 5.37; N, 18.65%. Found: C, 76.09; H, 5.40; N, 18.45%.

2-(3,5-Dinitrophenyl)-1*H*-benzimidazole as precursor **3**

Procedure for the synthesis of precursor **1** was repeated for the reaction of *o*-phenylenediamine with 3,5-dinitrophenylbenzoic acid to afford precursor **3** (90.07%), mp = 177–179 °C, colour = yellow. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 7.40–7.42 (d, J = 8.00 Hz, 1H, Ph–H), 7.44–7.46 (d, J = 8.00 Hz, 1H, Ph–H), 7.72–7.74 (t, J = 7.58 Hz, 1H, Ph–H), 7.94–7.96 (m, 1H, Ph–H), 8.64 (s, 2H, Ph–H), 8.85 (s, 1H, Ar–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 115.2 (2 × CH), 125.8, 128.7 (2 × CH), 129.4 (2 × CH), 134.8, 144.2 (2 × C), 150.7 (2 × C), 156.1 ppm. λ_{max} in nm (log ϵ_{max}): 218 (4.6522), 248 (4.6365). FT-IR (ν_{max} in cm^{-1}): 3349 (N–H), 3172 (C–H aromatic), 3101 (C–H aromatic), 1606 (C=C), 1572 (C=N), 1543 (NO_2 asym.), 1344 (NO_2 sym.). Anal. Calcd for $\text{C}_{13}\text{H}_8\text{N}_4\text{O}_4$ (284.22): C, 55.13; H, 2.49; N, 19.78%. Found: C, 54.98; H, 2.54; N, 19.92%.

Overall protocol towards accessing 1,2-disubstituted-1*H*-benzimidazole **4a-f**

Similar procedure for **2a-f** was repeated herein using 2-(3,5-dinitrophenyl)-1*H*-benzimidazole **3** as the precursor which reacted with substrates **a-f** to afford 1-substituted-2-(3,5-dinitrophenyl)-1*H*-benzimidazoles **4a-f**.

1-(2-(3,5-Dinitrophenyl)-1*H*-benzimidazole-1-yl)ethanone **4a**

Yield 72.58%, mp = 250–252 °C, colour = brown. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 2.59 (s, 3H, CH_3 -CO), 7.40–7.42 (d, J = 8.00 Hz, 1H, Ph–H), 7.44–7.46 (d, J = 8.00 Hz, 1H, Ph–H), 7.72–7.74 (t, J = 7.56 Hz, 1H, Ph–H), 7.94–7.96 (m, 1H, Ph–H), 8.64 (s, 2H, Ph–H), 8.85 (s, 1H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 28.9 (CH_3), 116.6 (2 × CH), 126.1, 128.9 (2 × CH), 129.7 (2 × CH), 135.0, 144.3 (2 × C), 150.5 (2 × C), 156.3, 174.1 (C=O) ppm. λ_{max} in nm (log ϵ_{max}): 218 (4.3729), 248 (4.3050). FT-IR (ν_{max} in cm^{-1}): 3106 (C–H aromatic), 2854 (C–H aliphatic), 1699 (C=O amide), 1622 (C=C), 1580 (C=N), 1541 (NO_2 asym.), 1346 (NO_2 sym.). Anal. Calcd for $\text{C}_{15}\text{H}_{10}\text{N}_4\text{O}_5$ (326.26): C, 55.22; H, 3.09; N, 17.17%. Found: C, 55.13; H, 2.89; N, 17.08%.

2-(3,5-Dinitrophenyl)-1-(phenylsulfonyl)-1*H*-benzimidazole **4b**

Yield 62.77%, mp > 300 °C, colour = gray. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 7.15–7.19 (m, 5H, Ph–H), 7.40–7.46 (m, 2H, Ph–H), 7.72–7.74 (t, J = 7.60 Hz, 1H, Ph–H), 7.94–7.96 (m, 1H, Ph–H), 8.64 (s, 2H, Ph–H), 8.85 (s, 1H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 116.6 (2 × CH), 119.6 (2 × CH), 126.1, 128.9 (2 × CH), 129.7 (2 × CH), 132.5, 135.0 (2 × C), 137.5 (2 × CH), 144.3 (2 × C), 150.5, 152.0, 156.3 ppm. λ_{max} in nm (log ϵ_{max}): 215 (3.9395), 251 (4.1492). FT-IR (ν_{max} in cm^{-1}): 3050 (C–H aromatic), 2855 (C–H aliphatic), 1620 (C=C), 1580 (C=N), 1541 (NO_2 asym.), 1376 (SO_2), 1346 (NO_2 sym.), 1185 (SO_2 2nd band). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{N}_4\text{O}_6\text{S}$ (434.39): C, 53.77; H, 2.85; N, 13.20%. Found: C, 53.95; H, 3.03; N, 13.31%.

2-(3,5-Dinitrophenyl)-1-tosyl-H-benzimidazole **4c**

Yield 98.77%, mp = 112 °C, colour = brown. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 2.60 (s, 3H, CH₃–Ar), 6.90–6.92 (d, *J* = 8.00 Hz, 2H, Ph–H), 7.15–7.17 (d, *J* = 8.00 Hz, 2H, Ph–H), 7.40–7.44 (m, 2H, Ph–H), 7.73–7.75 (t, *J* = 7.54 Hz, 1H, Ph–H), 7.93–7.95 (m, 1H, Ph–H), 8.60 (s, 2H, Ph–H), 8.84 (s, 1H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 21.7 (CH₃), 116.6 (2 × CH), 119.6 (2 × CH), 124.8, 156.3, 152.1, 150.5, 144.3 (2 × C), 137.5 (2 × CH), 135.0, 132.5, 129.5 (2 × CH), 126.0, 128.9 (2 × CH) ppm. λ_{max} in nm (log ε_{max}): 242 (4.6138). FT-IR (ν_{max} in cm^{−1}): 3105 (C–H aromatic), 2852 (C–H aliphatic), 1620 (C=C), 1575 (C=N), 1540 (NO₂ asym.), 1377 (SO₂), 1345 (NO₂ sym.). Anal. Calcd for C₂₀H₁₄N₄O₆S (438.41): C, 54.79; H, 3.22; N, 12.78%. Found: C, 54.62; H, 3.16; N, 12.94%.

1-(3-Chlorobenzyl)-2-(3,5-dinitrophenyl-1H-benzimidazole **4d**

Yield 71.16%, mp > 300 °C, colour = black. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 3.72 (s, 2H, CH₂–Ar), 7.11–7.14 (m, 1H, Ph–H), 7.25–7.27 (d, *J* = 7.56 Hz, 2H, Ph–H), 7.40–7.46 (m, 2H, Ph–H), 7.72–7.74 (t, *J* = 7.60 Hz, 1H, Ph–H), 7.94–7.96 (d, *J* = 7.22 Hz, 1H, Ph–H), 8.21 (s, 1H, Ph–H), 8.64 (s, 2H, Ph–H), 8.85 (s, 1H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 48.0 (CH₂), 115.2, 116.2 (2 × CH), 119.2, 125.1, 126.1, 128.9 (2 × CH), 129.7 (2 × CH), 130.0, 135.0, 139.1, 144.0 (2 × C), 150.4 (2 × C), 156.1 ppm. λ_{max} in nm (log ε_{max}): 212 (3.7076), 248 (4.1986), 467 (2.6020), 470 (2.6020). FT-IR (ν_{max} in cm^{−1}): 2924, 2854 (CH aliphatic), 1612 (C=C Aromatic), 1584 (C=N imine), 1501 (NO₂ asym.). Anal. Calcd for C₂₀H₁₃N₄O₄Cl (408.79): C, 58.76; H, 3.21; N, 13.71%. Found: C, 58.88; H, 3.32; N, 13.89%.

5-(2-(3,5-Dinitrophenyl)-1H-benzimidazol-1-yl)pyrimidine-2,4 (1H,3H)-dione **4e**

Yield 61.04%, mp > 300 °C, colour = gray. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 7.40–7.42 (t, *J* = 7.24 Hz, 1H, Ph–H), 7.44–7.46 (m, 1H, Ph–H), 7.72–7.74 (d, *J* = 8.00 Hz, 2H, Ph–H), 7.94–7.96 (s, 1H, Ph–H), 8.64 (s, 2H, Ph–H), 8.85 (s, 1H, Ph–H), 11.05 (s, 1H, NH), 11.55 (s, 1H, NH). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 116.6 (2 × CH), 126.1, 128.9 (2 × CH), 129.7 (2 × CH), 135.0, 138.7, 140.1, 144.3 (2 × C), 150.5 (2 × C), 156.3, 169.0 (C=O), 169.7 (C=O) ppm. λ_{max} in nm (log ε_{max}): 220 (4.0234), 242 (4.8730), 311 (4.0790). FT-IR (ν_{max} in cm^{−1}): 3362 (N–H), 3217 (N–H), 3050 (C–H aromatic), 1685 (C=O amide), 1612 (C=C aromatic), 1575 (C=N), 1536 (NO₂ asym.), 1344 (NO₂ sym.). Anal. Calcd for C₁₇H₁₀N₆O₆ (394.30): C, 51.78; H, 2.56; N, 21.31%. Found: C, 51.97; H, 2.69; N, 21.51%.

4-(2-(3,5-Dinitrophenyl)-1H-benzimidazole-1-yl)aniline **4f**

Yield 66.85%, mp > 300 °C, colour = gray. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 4.50 (s, 2H, NH₂), 6.90–6.92 (d, *J* = 8.00 Hz, 2H, Ph–H), 7.15–7.17 (d, *J* = 8.00 Hz, 2H, Ph–H), 7.40–7.44 (m, 2H, Ph–H), 7.73–7.76 (t, *J* = 7.84 Hz, 1H, Ph–H), 7.93–7.96 (m, 1H, Ph–H), 8.69 (s, 2H, Ph–H), 8.90 (s, 1H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 115.2, 116.6 (2 × CH), 118.3 (2 × CH), 125.3, 126.1, 128.9 (2 × CH), 129.7 (2 × CH), 135.0, 144.1 (2 × C), 147.0, 147.8, 150.5 (2 × C), 155.4 ppm. λ_{max} in nm (log ε_{max}): 235 (5.4130), 302 (4.8040). FT-IR (ν_{max} in cm^{−1}): 3472 (N–H), 3363 (N–H), 3214, 1620 (C=C aromatic), 1536 (NO₂ asym.), 1321, 1377 (NO₂sym.). Anal. Calcd for C₁₉H₁₃N₅O₄ (375.34): C, 60.80; H, 3.49; N, 18.66%. Found: C, 61.00; H, 3.68; N, 18.84%.

2-Benzyl-1H-benzimidazole as precursor **5**

Procedure for the synthesis of precursor **1** was repeated for the reaction of *o*-phenylenediamine with phenyl acetic acid to afford precursor **5** (79.12%), mp = 108–110 °C, colour = brown. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 3.56 (s, 2H, CH₂–Ar), 6.35–6.38 (dd, *J*₁ = 3.48 Hz, *J*₂ = 9.12 Hz, 1H, Ph–H), 6.48–6.50 (dd, *J*₁ = 3.48 Hz, *J*₂ = 8.00 Hz, 1H, Ph–H), 7.12 (s, 5H, Ph–H), 7.24–7.26 (d, *J* = 9.12 Hz, 1H, Ph–H), 7.29–7.31 (d, *J* = 8.00 Hz, 1H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 34.9 (CH₂), 115.2 (2 × CH), 123.4 (2 × CH), 125.8, 128.7 (2 × CH), 129.4 (2 × CH), 136.9, 138.8 (2 × C), 142.7 ppm. λ_{max} in nm (log ε_{max}): 257 (5.4181), 275 (5.4133), 314 (4.4842). FT-IR (ν_{max} in cm^{−1}): 3415 (N–H), 2924 (C–H aliphatic), 2854 (C–H aliphatic), 1638 (C=C), 1587 (C=N imine). Anal. Calcd for C₁₄H₁₂N₂ (208.26): C, 80.74; H, 5.81; N, 13.45%. Found: C, 80.80; H, 6.01; N, 13.44%.

Overall protocol towards accessing 1,2-disubstituted-1H-benzimidazole **6a-f**

Similar procedure for the synthesis of **2a-f** was repeated herein using 2-benzyl-1H-benzimidazole **5** as the precursor which reacted with substrates **a-f** to afford 1-substituted-2-benzyl-1H-benzimidazoles **6a-f**.

1-(2-Benzyl-1H-benzimidazole-1-yl)ethanone **6a**

Yield 90.11%, mp = N.D. (Oily), colour = black. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 2.67 (s, 3H, CH₃CO), 3.56 (s, 2H, CH₂–Ar), 6.36–6.39 (dd, *J*₁ = 3.48 Hz, *J*₂ = 9.12 Hz, 1H, Ph–H), 6.48–6.50 (dd, *J*₁ = 3.48 Hz, *J*₂ = 8.00 Hz, 1H, Ph–H), 7.09 (s, 5H, Ph–H), 7.24–7.26 (d, *J* = 9.12 Hz, 1H, Ph–H), 7.29–7.31 (d, *J* = 8.00 Hz, 1H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 28.9 (CH₃), 34.9 (CH₂), 115.2 (2 × CH), 123.5 (2 × CH), 125.9, 128.7 (2 × CH), 129.5 (2 × CH), 136.9, 138.9 (2 × C), 142.8 ppm. λ_{max} in nm (log ε_{max}): 227 (2.2300), 315 (2.5057). FT-IR (ν_{max} in cm^{−1}): 2922 (C–H aliphatic), 2855 (C–H aliphatic), 1685 (C=O), 1620 (C=C), 1575 (C=N imine). Anal. Calcd for C₁₆H₁₄N₂O (250.30): C, 76.78; H, 5.64; N, 11.19%. Found: C, 76.94; H, 5.59; N, 11.00%.

2-Benzyl-1-(phenylsulfonyl)-1H-benzimidazole **6b**

Yield 87.48%, mp = N.D. (Oily), colour = black. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 3.56 (s, 2H, CH₂–Ar), 6.35–6.38 (dd, *J*₁ = 3.60 Hz, *J*₂ = 9.26 Hz, 1H, Ph–H), 6.48–6.50 (dd, *J*₁ = 3.60 Hz, *J*₂ = 8.00 Hz, 1H, Ar–H), 6.91–6.95 (m, 3H, Ph–H), 7.12 (s, 5H, Ph–H), 7.13–7.17 (m, 3H, Ph–H), 7.22–7.24 (d, *J* = 9.26 Hz, 1H, Ph–H), 7.30–7.32 (d, *J* = 8.00 Hz, 1H, Ph–H), 7.44–7.46 (d, *J* = 8.66 Hz, 2H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 34.7 (CH₂), 115.2 (2 × CH), 123.4 (2 × CH), 125.8, 128.1 (2 × CH), 128.7 (2 × CH), 129.0 (2 × CH), 129.9 (2 × CH), 133.9, 136.8, 137.9, 138.7 (2 × C), 142.9 ppm. λ_{max} in nm (log ε_{max}): 215 (4.4669), 254 (4.4540). FT-IR (ν_{max} in cm^{−1}): 2922, 2850 (CH aliphatic), 1620 (C=C aromatic), 1575 (C=N imine), 1375 (SO₂), 1185 (SO₂ 2nd band). Anal. Calcd for C₂₀H₁₆N₂O₂S (348.42): C, 68.94; H, 4.63; N, 8.04%. Found: C, 69.00; H, 4.82; N, 7.89%.

2-Benzyl-1-tosyl-1H-benzimidazole **6c**

Yield 93.25%, mp = N.D. (Oily), colour = brown. ¹H NMR (400 MHz, DMSO-*d*₆) δ_H: 2.76 (s, 3H, CH₃–Ar), 3.56 (s, 2H, CH₂–Ar), 6.35–6.38 (dd, *J*₁ = 3.60 Hz, *J*₂ = 9.26 Hz, 1H, Ph–H), 6.48–6.49 (dd, *J*₁ = 3.60 Hz, *J*₂ = 8.00 Hz, 1H, Ph–H), 6.92–6.94 (d, *J* = 8.76 Hz, 2H, Ph–H), 7.12 (s, 5H, Ph–H), 7.22–7.24 (d, *J* = 9.26 Hz, 1H, Ph–H), 7.30–7.32 (d, *J* = 8.00 Hz, 1H, Ph–H), 7.44–7.46 (d, *J* = 8.76 Hz, 2H, Ph–H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ_C: 21.5 (CH₃), 34.9 (CH₂),

115.2 (2 × CH), 123.4 (2 × CH), 125.8, 128.1, (2 × CH), 128.7 (2 × CH), 129.5 (2 × CH), 130.1 (2 × CH), 134.7, 136.7, 138.9 (2 × C), 139.6, 142.6 ppm. λ_{\max} in nm (log ϵ_{\max}): 233 (5.3679), 299 (4.7896). FT-IR (ν_{\max} in cm^{-1}): 2924, 2854 (CH aliphatic), 1648 (C=C aromatic), 1562 (C=N imine), 1376 (SO_2), 1185 (SO_2 2nd band). Anal. Calcd for $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_2\text{S}$ (362.44): C, 69.59; H, 5.01; N, 7.73%. Found: C, 69.51; H, 4.88; N, 7.82%.

2-Benzyl-1-(3-chlorobenzyl)-1H-benzimidazole **6d**

Yield 91.83%, mp = N.D. (Oily), colour = brown. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 3.42 (s, 2H, $\text{CH}_2\text{-Ar}$), 3.58 (s, 2H, $\text{CH}_2\text{-Ar}$), 6.35–6.38 (dd, $J_1 = 3.48$ Hz, $J_2 = 9.18$ Hz, 1H, Ph–H), 6.48–6.50 (dd, $J_{\text{m1}} = 3.48$ Hz, $J_2 = 8.00$ Hz, 1H, Ph–H), 7.10 (s, 5H, Ph–H), 7.17–7.19 (dd, $J_1 = 7.20$ Hz, $J_2 = 7.82$ Hz, 1H, Ph–H), 7.23–7.25 (d, $J = 9.18$ Hz, 1H, Ph–H), 7.31–7.33 (d, $J = 8.00$ Hz, 1H, Ph–H), 7.37–7.38 (d, $J = 7.82$ Hz, 1H, Ph–H), 7.51–7.52 (d, $J = 7.20$ Hz, 1H, Ph–H), 8.21 (s, 1H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 33.8 (CH_2), 48.0 (CH_2), 115.2 (2 × CH), 123.4 (2 × CH), 125.1, 125.6, 125.8, 128.7 (2 × CH), 128.9, 129.5 (2 × CH), 130.2, 134.4, 136.7, 137.8, 138.9 (2 × C), 142.5 ppm. λ_{\max} in nm (log ϵ_{\max}): 233 (5.3877), 251 (5.0266). FT-IR (ν_{\max} in cm^{-1}): 2984 (C–H aliphatic), 1646 (C=C), 1565 (C=N), 652 (C–Cl). Anal. Calcd for $\text{C}_{20}\text{H}_{17}\text{ClN}_2$ (332.83): C, 75.78; H, 5.15; N, 8.42%. Found: C, 75.70; H, 4.01; N, 8.25%.

5-(2-Benzyl-1H-benzimidazole-1-yl)pyrimidine-2,4(1H,3H)-dione **6e**

Yield 80.58%, mp = N.D. (Oily), colour = black. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 3.56 (s, 2H, $\text{CH}_2\text{-Ar}$), 6.34–6.37 (dd, $J_1 = 3.44$ Hz, $J_2 = 9.18$ Hz, 1H, Ph–H), 6.48–6.50 (dd, $J_1 = 3.44$ Hz, $J_2 = 8.00$ Hz, 1H, Ph–H), 7.11 (s, 5H, Ph–H), 7.23–7.25 (d, $J = 9.18$ Hz, 1H, Ph–H), 7.30–7.32 (d, $J = 8.00$ Hz, 1H, Ph–H), 7.95–7.97 (s, 1H, Ph–H), 11.05 (s, 1H, NH), 11.55 (s, 1H, NH). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 34.8 (CH_2), 115.2 (2 × CH), 119.9, 123.3 (2 × CH), 125.7, 128.7 (2 × CH), 129.4 (2 × CH), 136.7, 138.5 (2 × C), 142.9, 150.0, 169.2 (C=O), 169.7 (C=O) ppm. λ_{\max} in nm (log ϵ_{\max}): 218 (4.6532), 248 (4.5775), 293 (4.1399), 437 (3.4914). FT-IR (ν_{\max} in cm^{-1}): 3360 (N–H), 3217 (N–H), 3050 (C–H aromatic), 1685 (C=O amide), 1615 (C=C aromatic), 1575 (C=N). Anal. Calcd for $\text{C}_{18}\text{H}_{14}\text{N}_4\text{O}_2$ (318.33): C, 67.91; H, 4.43; N, 17.60%. Found: C, 68.10; H, 4.57; N, 17.77%.

4-(2-Benzyl-1H-benzimidazole-1-yl) aniline **6f**

Yield 86.24%, mp = N.D. (Oily), colour = brown. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ_{H} : 3.56 (s, 2H, $\text{CH}_2\text{-Ar}$), 4.50 (s-br, 2H, NH_2), 6.35–6.38 (dd, $J_1 = 3.46$ Hz, $J_2 = 9.12$ Hz, 1H, Ph–H), 6.48–6.50 (dd, $J_1 = 3.46$ Hz, $J_2 = 8.00$ Hz, 1H, Ph–H), 7.11 (s, 5H, Ph–H), 7.16–7.18 (d, $J = 8.20$ Hz, 2H, Ph–H), 7.24–7.26 (d, $J = 9.12$ Hz, 1H, Ph–H), 7.29–7.31 (d, $J = 8.00$ Hz, 1H, Ph–H), 7.44–7.46 (d, $J = 8.20$ Hz, 2H, Ph–H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ_{C} : 34.9 (CH_2), 115.2 (2 × CH), 118.7 (2 × CH), 123.4 (2 × CH), 125.8, 126.2 (2 × CH), 128.7 (2 × CH), 129.4 (2 × CH), 136.9, 138.8 (2 × C), 142.7, 146.8, 148.7 ppm. λ_{\max} in nm (log ϵ_{\max}): 230 (5.2753), 251 (5.1322), 299 (4.8401). FT-IR (ν_{\max} in cm^{-1}): 3404 (N–H interfered by OH), 2985 (C–H aliphatic), 1610 (C=C aromatic), 1572 (C=N). Anal. Calcd for $\text{C}_{20}\text{H}_{17}\text{N}_3$ (299.37): C, 80.24; H, 5.72; N, 14.04%. Found: C, 80.21; H, 5.90; N, 13.85%.

Antibacterial sensitivity testing of compounds **1–6f**

All the synthesized compounds (**1–6f**) and gentamicin were screened for antibacterial activity on four bacterial strains using agar well diffusion method [26]. The detailed was as attached in Supplementary Materials.


Minimum inhibitory/bactericidal concentration (MIC and MBC) testing

The minimum inhibitory concentration (MIC) test on the chosen organisms was carried out via serial dilution technique [27] and the concentration range was from 500.00 to 15.63 $\mu\text{g/mL}$, while the minimum bactericidal concentration (MBC) was determined by a standard method [26]. The detailed description of the procedures for the determination of MIC and MBC were as presented in the supplementary material.


Results and discussion

Based on the enthusiastic outcome of an extensive review on functionalized benzimidazole [3] and in the continuation of the research effort in the area of benzo-fused imidazole moieties [6,28], the synthesis of functionalized 2-substituted and 1,2-disubstituted benzimidazole derivatives was herein reported to evaluate their antibacterial activities. Although, various derivatives of benzimidazole moieties have been synthesized and reported in literature; high temperature for the reflux process and the uses harsh reaction condition in the presence of strong and concentrated acids such as HCl [4,6] have been involved. On the contrary, the synthesis of 2-(1H-benzimidazol-2-yl)aniline precursor **1**, was achieved herein by the use of eco-friendly reaction of *o*-phenylenediamine with anthranilic acid using a catalytic amount of NH_4Cl in the presence of ethyl alcohol as a solvent at refluxing temperature of 60–70 °C (Scheme 1a). The synthetic modification of NH_2 functional side chain of the reactive intermediate **1** was conducted by reacting it with six different electrophile-releasing substrates to furnish **2a–f**. Prior to this, the reaction optimization study was carried out using two main parameters. First, solvent dependent condition was investigated using the synthesis of *N*-(2-(1H-benzimidazole-2-yl)phenyl)acetamide **2a** via the reaction of **1** with acetyl chloride in either tetrahydrofuran (THF), ethanol or acetonitrile at room temperature for 24 h; this gave yield of 87.7%, 30%, and 25% respectively. In addition, the thermodynamic dependent kinetic of the synthesis of **2a** was evaluated using the comparative study of the synthesis in THF at room temperature, 60 °C, and 120 °C. It was unveiled that room temperature gave the highest yield (87.71%) followed by 60 °C (38.24%), while there was no isolated product at 120 °C. Thus, the same reaction condition was adopted for reaction of **1** with the remaining five electrophile-releasing substrates **b–f** to produce diverse functionalized 2-substitutedbenzimidazole derivatives **2b–f** (Scheme 1b). According to another route [29], the *o*-phenylenediamine reacted with 3,5-dinitrobenzoic acid to achieve **3** Scheme 2a, which was subsequently treated with the earlier reported six electrophile-releasing substrates to furnish 1,2-disubstituted benzimidazole derivatives **4a–f** (Scheme 2b) in varying yields. Finally, the last precursor **5** in this present work was prepared by the condensation of *o*-phenylenediamine with phenyl acetic acid (Scheme 3a). Precursor **5** was eventually reacted with the six electrophile-releasing substrates to access the 1,2-disubstituted benzimidazole derivatives **6a–f** (Scheme 3b).

Physicochemical parameter data were presented in experimental section alongside the elemental analysis data. The result of elemental analysis for the % calculated agreed with % found for C, H N of all the synthesized compounds with high state of accuracy (the difference was not more than ± 0.20 in all cases). Furthermore, the spectroscopic characterization of the targeted templates was carried out using IR and UV, ^1H , and ^{13}C NMR analysis. The ^1H NMR spectra of the compounds were run in $\text{DMSO}-d_6$ at 400 MHz with chemical shift values recorded in ppm. The aryl-linked CH_3 of **2c**, **4c**, and **6c** resonated upfield at δ 2.60–2.80 ppm as 3H singlets and acetyl-linked CH_3 of **2a**, **4a**, and **6a** were seen upfield as 3H sin-




Scheme 1. (a) Synthesis of the precursor **1** (b) Synthesis of 2-substitutedbenzimidazoles **2a-f**.



Scheme 2. (a) Synthesis of the precursor **3** (b) Synthesis of 1,2-disubstitutedbenzimidazoles **4a-f**.

glets at δ 2.60–2.67 ppm. The aryl-linked CH_2 of **2d**, **4d**, and **5** as well as **6a-f** were noticed as singlets at δ 3.42–3.73 ppm. Signals of all aryl H appeared downfield of TMS around δ 6.31–8.90 ppm, NH_2 of amine in **1**, **2f**, **4f**, and **6f** were broad singlets at δ 4.50–5.32 ppm. The most downfield signals were that of N–H of amide

found as singlets in compounds **2e**, **4e**, and **6e** at δ 11.02–11.56 ppm. The ^{13}C NMR spectra were run in $\text{DMSO}-d_6$ at 100 MHz with chemical shift values recorded in ppm. On the overall, the ^{13}C NMR spectra of the structure-based benzimidazole derivatives varied from 21.3 ppm for CH_3 of compound **2c** to



The general sensitivity testing was evaluated using the *in vitro* screening of the synthesized compounds against four bacterial isolates (*Staphylococcus aureus*, *Bacillus licheniformis*, *Proteus vulgaris*, and *Pseudomonas aeruginosa*). Gentamicin was used as the positive control in this study. The justification of gentamicin as a clinical standard was due to the mode of action, which involved irreversible binding at ribosomal level, thereby signaling to obstruct

and interrupt protein synthesis [31]. Agar diffusion method was used for the sensitivity testing and the diameters of zones of inhibition (Z. O. I) were documented in millimeter (Table 1). Although, large zones of inhibition were noticed for most of the targeted benzimidazole derivatives final products against the screened organisms, resistance was observed in few cases such as **6c** against *S. aureus*; **4d-f**, **6a** against *Bacillus licheniformis*; **2a**, **2c**, **4a**, **4d**, and **6f** against *Proteus vulgaris*; **2e**, **4d-f**, and **6a** against *Pseudomonas aeruginosa*, and **6f** developed resistant against the effect of gentamicin. Overall, the largest zone of inhibition (40.00 ± 0.10 mm) was recorded for **1** against *S. aureus*, while the lowest zone of inhibition (15.00 ± 0.08 mm) was recorded for **3** against *S. aureus*. In comparison with gentamicin, all synthesized compounds, except **3**, had better activity with higher zones of inhibition against growth potential of *S. aureus*. This means that the array of compounds synthesized herein might be a possible replacement for gentamicin on infectious disease caused by the *S. aureus* or enhance the potency of gentamicin where resistance issues occur. Compared to gentamicin, all compounds except **3** and **6c** showed larger zones of inhibition against growth of *S. aureus* (i.e. >16 mm); and except **4d-f** and **6c** against *B. licheniformis* (i.e. >16 mm), while **3** and **4c** exhibited approximately the same Z.O.I. as gentamicin against *S. aureus* (15 mm) and *B. licheniformis* (16 mm), respectively. Interestingly, more than 75% of the targeted benzimidazole derivatives were active on *P. vulgaris* with large zones of inhibition, whereas this organism was resistant to gentamicin (Table 1). Similarly, more than 75% of the targeted benzimidazole derivatives (Z.O. I = 25.00 ± 0.08 to 38.00 ± 0.12 mm) were more active than gentamicin (20.00 ± 0.08 mm) upon *P. aeruginosa*. In the present study, the choice of *S. aureus* and *E. coli* was due to the broad array of pathogenic infections and precarious health issue that are associated with these bacterial strains [32] and wide reported occurrence of resistant strain of *S. aureus* [1]. *S. aureus* has been reported to have strong relationship with death rate increase in human population, prolong admission of patients in hospitals, and the infections caused by this bacterial strain are expensive to treat [33].

Table 1
Antibacterial sensitivity testing with zones of inhibition in millimetre.

Compound No.	<i>S. aureus</i>	<i>B. licheniformis</i>	<i>P. vulgaris</i>	<i>P. aeruginosa</i>
1	40.00 ± 0.10	26.00 ± 0.08	35.00 ± 0.09	38.00 ± 0.12
2a	27.00 ± 0.09	26.00 ± 0.08	R	28.00 ± 0.08
2b	30.00 ± 0.09	20.00 ± 0.08	25.00 ± 0.09	30.00 ± 0.08
2c	35.00 ± 0.09	25.00 ± 0.08	R	28.00 ± 0.08
2d	30.00 ± 0.10	28.00 ± 0.09	30.00 ± 0.09	30.00 ± 0.09
2e	32.00 ± 0.12	26.00 ± 0.08	35.00 ± 0.08	R
2f	30.00 ± 0.11	30.00 ± 0.12	28.00 ± 0.08	30.00 ± 0.08
3	15.00 ± 0.08	20.00 ± 0.08	30.00 ± 0.12	28.00 ± 0.09
4a	20.00 ± 0.10	24.00 ± 0.09	R	25.00 ± 0.08
4b	30.00 ± 0.10	18.00 ± 0.08	20.00 ± 0.08	25.00 ± 0.08
4c	25.00 ± 0.10	16.00 ± 0.08	20.00 ± 0.08	25.00 ± 0.08
4d	30.00 ± 0.09	R	R	R
4e	30.00 ± 0.10	R	20.00 ± 0.08	R
4f	30.00 ± 0.12	R	23.00 ± 0.08	R
5	30.00 ± 0.09	28.00 ± 0.09	32.00 ± 0.12	38.00 ± 0.08
6a	19.00 ± 0.08	R	18.00 ± 0.08	R
6b	35.00 ± 0.09	22.00 ± 0.08	30.00 ± 0.10	28.00 ± 0.08
6c	R	18.00 ± 0.08	32.00 ± 0.08	26.00 ± 0.08
6d	32.00 ± 0.10	26.00 ± 0.08	35.00 ± 0.12	30.00 ± 0.08
6e	25.00 ± 0.08	24.00 ± 0.08	25.00 ± 0.08	30.00 ± 0.09
6f	20.00 ± 0.08	26.00 ± 0.08	R	28.00 ± 0.08
Gentamicin	16.00 ± 0.09	16.00 ± 0.08	R	20.00 ± 0.08

R = Resistance. Mean ± SD of triplicate determination.

Due to heat stable toxin production, *S. aureus* is enlisted as one of the highly invasive organisms referred to as pyogenic cocci involved in numerous adverse infectious conditions in humans [28,32,34]. In view of the reported predicament aforementioned and large zones of inhibition experienced via action of the benzimidazole framework herein on *S. aureus*, motivation was enhanced to investigate the activity index (A.I.) of the synthesized compounds against this organism (Fig. 1). It is quite impressive to note that all the synthesized benzimidazole motifs herein, showed better activity indices than gentamicin against *S. aureus* except **3** and **6c**. The compound **3** competed favourably with gentamicin (A. I. \approx 1.00) while **6c** developed resistance; hence could not have activity index.

Furthermore, based on high susceptibility of the organisms to the synthesized benzimidazole templates 1–6f and broad spectrum of activity observed herein, the MIC testing was carried out to determine the lowest concentration of the compound solution that conveniently inhibited the bacterial growth (Table 2). It was carried out using serial dilution method (500, 250, 125, 62.50, 31.25, and 15.63 μ g/mL) via an earlier reported method [27]. The MIC of benzimidazole derivatives upon *S. aureus* varied from 15.63 \pm 1.63 to 250 \pm 2.66 μ g/mL; against *B. licheniformis* varied

from 62.50 \pm 2.04 to 250 \pm 2.65 μ g/mL; against *P. vulgaris* varied from 31.25 \pm 1.94 to 250 \pm 2.65 μ g/mL; and against *P. aeruginosa* ranged from 15.63 \pm 1.63 to 125 \pm 2.45 μ g/mL. The best activity against *S. aureus* was observed in the precursor **1** among the series of the 2-substituted benzimidazoles **1–2f**. The presence of amino functionality in **1** and the ready availability of its lone pair of electron for coordination led to improved activity. The activity decreases after the NH₂ had underwent substitution as contained in **2a–f**, except for **2c** and **2e**, which competed favorably with **1**. This means that incorporation of *p*-toluenesulfonamido (in **2c**) and pyrimidinedione (in **2e**) played significant role as essential pharmacophores in increased activity observed in **2c** and **2e** against *S. aureus*. On the contrary, the series of 1,2-disubstituted benzimidazoles **4a–f** were more active than the precursor **3** from which they were derived. This showed that additional substitution on position 1 of compound **3** to afford 1,2-disubstitution in **4a–f**, was a worthwhile adventure in increasing the bioactivity of precursor **3**, since the series of 1,2-disubstituted benzimidazole products **4a–f** resulted in drastic improvement of growth inhibition in *S. aureus*. In addition to MIC testing, the minimum bactericidal concentration (MBC) testing was determined to authenticate the lowest concentration of the benzimidazole solution that causes death




Fig. 1. Comparative study of the activity index of synthesized benzimidazoles and gentamicin.

Table 2Minimum inhibitory concentration (MIC) in $\mu\text{g/mL}$ of targeted benzimidazoles, **1–6f**.

Compound No↓	<i>S. aureus</i>	<i>B. licheniformis</i>	<i>P. vulgaris</i>	<i>P. aeruginosa</i>
1	15.63 \pm 1.64	125.00 \pm 2.44	62.50 \pm 2.04	31.25 \pm 1.98
2a	31.25 \pm 1.98	125.00 \pm 2.43	N.D.	62.50 \pm 2.01
2b	31.25 \pm 1.97	250.00 \pm 2.67	125.00 \pm 2.44	31.25 \pm 1.94
2c	15.63 \pm 1.67	125.00 \pm 2.44	N.D.	62.50 \pm 2.03
2d	31.25 \pm 1.96	125.00 \pm 2.44	62.50 \pm 2.01	62.50 \pm 2.03
2e	15.63 \pm 1.63	250.00 \pm 2.67	62.50 \pm 2.01	N.D.
2f	31.25 \pm 1.94	62.50 \pm 2.04	125.00 \pm 2.44	31.25 \pm 1.94
3	250.00 \pm 2.66	250.00 \pm 2.67	125.00 \pm 2.44	62.50 \pm 2.01
4a	125.00 \pm 2.45	125.00 \pm 2.44	N.D.	62.50 \pm 2.02
4b	31.25 \pm 1.95	250.00 \pm 2.66	250.00 \pm 2.67	62.50 \pm 2.01
4c	62.50 \pm 2.01	250.00 \pm 2.66	250.00 \pm 2.66	31.25 \pm 1.97
4d	31.25 \pm 1.98	N.D.	N.D.	N.D.
4e	15.63 \pm 1.65	N.D.	250.00 \pm 2.65	N.D.
4f	31.25 \pm 1.94	N.D.	250.00 \pm 2.67	N.D.
5	15.63 \pm 1.67	62.50 \pm 2.04	62.50 \pm 2.03	15.63 \pm 1.63
6a	250.00 \pm 2.66	N.D.	125.00 \pm 2.44	N.D.
6b	15.63 \pm 1.63	250.00 \pm 2.65	62.50 \pm 2.04	62.50 \pm 2.02
6c	N.D.	250.00 \pm 2.67	31.25 \pm 1.94	125.00 \pm 2.45
6d	31.25 \pm 1.94	125.00 \pm 2.44	31.25 \pm 1.95	31.25 \pm 1.96
6e	125.00 \pm 2.45	125.00 \pm 2.45	125.00 \pm 2.44	62.50 \pm 2.02
6f	125.00 \pm 2.43	125.00 \pm 2.45	N.D.	62.50 \pm 2.01
Gentamicin	5.00 \pm 1.23	2.5 \pm 1.09	N.D.	7.5 \pm 1.45


N.D. = Not Determined. Mean \pm SD of triplicate determination.

of the bacterium targeted per time and the results are shown in Fig. 2. Apart from where the MBC was not determined (N.D.) due to occurrence of resistance, all other cases showed that the MBC values were double the MIC in each consideration, except in **6a** alone where MBC (500 $\mu\text{g/mL}$) was 4 times that of MIC (125 $\mu\text{g/mL}$) against *P. vulgaris*. From all indications, the lowest MBC trend was observed for the bio-assay screening of benzimidazole derivatives upon *S. aureus*, whereas the highest MBC was reported against *P. aeruginosa* as shown in Fig. 2.

Structure activity relationship (SAR) study

From the overview of the structure activity relationship (SAR) study, it was found that the nature of substituent on 1-position and 2-positions of the benzimidazole nucleus had significant effect on the antibacterial activity of the entire structures. The compounds series **2a–f** were structurally related in the core pharmacophoric 2-phenylbenzimidazole; the SAR study showed their

activity against *S. aureus* to be in the order **2c** \approx **2e** $>$ **2a** \approx **2b** \approx **2d** \approx **2f**. This means that the presence of electron donating CH_3 on *p*-toluenesulfonamide moieties in the 2-position of anilino side chain played a significant role in the improvement of activity, since its counterpart **2b** without CH_3 was far less active and stayed in the categories of **2a**, **2d**, and **2f** in its activity upon *S. aureus* growth inhibition. Considering **4a–f** on *S. aureus*, the activity varied in the order of: **4e** $>$ **4b** \approx **4d** \approx **4f** $>$ **4c** $>$ **4a**. Thus, electron withdrawing ability and π - π stacking character in pyrimidine-dione at 1-position of 2-(3,5-dinitrophenyl)benzimidazole core worked synergistically with the electron withdrawing NO_2 on benzene at 2-position to increase the activity of **4e**, thereby causing it to exhibit outstanding activity against *S. aureus* among the **4a–f** series. Based on the *in vitro* screening of **6a–f** against *S. aureus*, the order of activity was **6b** $>$ **6d** $>$ **6e** \approx **6f** $>$ **6a** $>$ **6c**. Hence, presence of benzenesulfonamido group on 1-position of 2-benzylbenzimidazole in series **6a–f** played a crucial role in activity boosting, making **6b** to be the most active among the group and more active as compared to its isomor-

**Fig. 2.** Graphical representation of minimum bactericidal concentration (MBC).

phic template **6c** where no activity was noticed. It was interesting to note that *p*-methyl group in **6c** which was the only group absent in **6b** provided the framework **6a–f** with antagonistic effect thereby causing total activity loss in **6c** as compared to **6b** against *S. aureus*.

Conclusions

Benzimidazole is an essential heterocyclic framework in agrochemicals, pharmaceuticals, and medicinal chemistry research. NH₄Cl catalyzed strategy was found to be efficient approach for accessing the reported benzimidazole precursor in good yield. Thus, mono- and disubstituted benzimidazole derivatives with improved medicinal potential were successfully synthesized via an elegant pathway. The findings of the *in vitro* screening unveiled the broad spectrum of activity of the synthesized benzimidazole templates in the present study. Among the series, the highest potency was exerted and experienced in 2-(1*H*-benzimidazol-2-yl)aniline, **1** and 2-benzyl-1-(phenylsulfonyl)-1*H*-benzimidazole, **6b**. It will be a worthwhile adventure to advance the work further for more guidance on the pharmacokinetic and pharmacodynamic study to ascertain probable candidature of the templates for future drug design.

Conflict of interest

The authors have declared no conflict of interest.

Compliance with Ethics Requirements

This article does not contain any studies with human or animal subjects.

Acknowledgements

OOA is thankful to The World Academy of Sciences for the sponsorship of this project under the TWAS Research Grants Programme in Basic Sciences for Individual Scientists (Grant No. 14-069 RG/CHE/AF/AC_1). Covenant University is also gratefully acknowledged for support for this present work.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jare.2017.09.003>.

References

- [1] Ajani OO, Familoni OB, Aderohunmu DV, Ogunniran KO, Adekoya JA, Olanrewaju IO. Comparative study of the antibacterial activity of *N,N*-diethylamido substituted *p*-toluenesulfonamides to their α -toluenesulfonamide counterparts. *Pak J Biol Sci* 2015;18:166–72.
- [2] Martins P, Jesus J, Santos S, Raposo LR, Roma-Rodrigues C, Baptista PV, et al. Heterocyclic anticancer compounds: recent advances and the paradigm shift towards the use of nanomedicine's tool box. *Molecules* 2015;20:16852–91.
- [3] Ajani OO, Aderohunmu DV, Ikpo CO, Adedapo EA, Olanrewaju IO. Functionalized benzimidazole scaffolds: privileged heterocycle for drug design in therapeutic medicine. *Arch Pharm* 2016;349:1–32.
- [4] Bahrami K, Khodaei MM, Kaviani I. A simple and efficient one-pot synthesis of 2-substituted benzimidazoles. *Synthesis* 2007;4:547–50.
- [5] Bahrami K, Khodaei MM, Naali F. Mild and highly efficient method for the synthesis of 2-arylbenzimidazoles and 2-arylbenzothiazoles. *J Org Chem* 2008;73:6835–7.
- [6] Ajani OO, Ezeoke EK, Edobor-Osoh A, Ajani OA. Facile synthesis and characterization of new 2,3-disubstituted benzimidazole derivatives. *Int Res J Pure Appl Chem* 2013;3:10–21.
- [7] Bahrami K, Khodaei MM, Naali F. H₂O₂/Fe(NO₃)₃-promoted synthesis of 2-aryl benzimidazoles and 2-arylbenzothiazoles. *Synlett* 2009;4:569–72.
- [8] Zhu CJ, Wei YY. An inorganic iodine-catalyzed oxidative system for the synthesis of benzimidazoles using hydrogen peroxide under ambient conditions. *Chemsuschem* 2011;4:1082–6.
- [9] Goswami B, Singh AK. Pharmacological activities of benzimidazole derivatives – Overview. *Int J Sci Inno Discov* 2012;2(1):121–36.
- [10] Kondaparla S, Agarwal P, Srivastava K, Puri SK, Katti SB. Design, synthesis and *in vitro* antiparasitic activity of some bisquinolines against chloroquine-resistant strain. *Chem Biol Drug Des* 2017;89(6):901–6.
- [11] Patil SA, Patil SA, Patil R. Medicinal application of (benz)imidazole- and indole-based macrocycles. *Chem Biol Drug Des* 2017;89:639–49.
- [12] El-Gohary NS, Shaaban MI. Synthesis and biological evaluation of a new series of benzimidazole derivatives as antimicrobial, antitumor-sensing and antitumor agents. *Eur J Med Chem* 2017;131:255–62.
- [13] Alasmay FAS, Snelling AM, Zain ME, Alafeefy AM, Awaad AS, Karodia N. Synthesis and evaluation of selected benzimidazole derivatives as potential antimicrobial agents. *Molecules* 2015;20:15206–23.
- [14] Mavrova AT, Yancheva D, Anastassova N, Anichina K, Zvezdanovic J, Djordjevic A, et al. Synthesis, electronic properties, antioxidant and antibacterial activity of some new benzimidazole. *Bioorg Med Chem* 2015;23(19):6317–26.
- [15] Shaharyar M, Mazumder A, Salahuddin, Garg R, Pandey RD. Synthesis, characterization and pharmacological screening of novel benzimidazole derivatives. *Arab J Chem* 2016;9: S342–S47.
- [16] Tang Y, Zhang F, Hu S, Cao Z, Wu Z, Jing W. Novel benzimidazole derivatives as corrosion inhibitors of mild steel in the acidic media. Part I: gravimetric, electrochemical, SEM and XPS studies. *Corros Sci* 2013;74:271–82.
- [17] Abdel-Maksoud M, El-Shokry M, Ismail G, Hafez S, El-Kholy A, Attia E, et al. Methicillin-resistant *Staphylococcus aureus* recovered from healthcare- and community-associated infections in Egypt. *Int J Bacteriol* 2016;2016:5.
- [18] Alotaibi FE, Bukhari EE. Emergence of Vancomycin-resistant *Enterococci* at a teaching hospital, Saudi Arabia. *Chin Med J (Engl)* 2017;130(3):340–6.
- [19] Davin-Regli A, Pagès JM. *Enterobacter aerogenes* and *Enterobacter cloacae*: versatile bacterial pathogens confronting antibiotic treatment. *Front Microbiol* 2015;6:392.
- [20] Mwanjika GO, Subbiah M, Buza J, Rugumisa BT, Call DR. A systematic review of antibiotic-resistant *Escherichia coli* and *Salmonella* data obtained from Tanzanian healthcare settings (2004–2014). *Afr J Microbiol Res* 2017;11(2):45–54.
- [21] Mwanza S, Joshi S, Nambozi M, Chileshe J, Malunga P, Kabuya JBB, et al. The return of chloroquine-susceptible *Plasmodium falciparum* malaria in Zambia. *Malar J* 2016;15:584. doi: <https://doi.org/10.1186/s12936-016-1637-3>.
- [22] Ayandiran TA, Dahunsi SO. Microbial evaluation and occurrence of multidrug multi-resistant organisms among the indigenous *Clarias* species in River Oluwa, Nigeria. *J King Saud Univ Sci* 2017;29:96–105.
- [23] More D. Sulpha medication allergies in: sulpha drug allergy. Available online at: <<http://allergies.about.com/od/medicationallergies/a/sulfa.htm>> [accessed on 1st of June, 2016].
- [24] Routledge PA, O'Mahony MS, Woodhouse KW. Adverse drug reactions in elderly patients. *Br J Clin Pharmacol* 2004;57(2):121–6.
- [25] Aslan MM, Westerhaus M, Herce M, Nakashima K, Farmer PE. Poverty, global health and infectious disease: lessons from Haiti and Rwanda. *Infect Dis Clin North Am* 2011;25(3):611–22.
- [26] Ajani OO, Nwinyi OC. Microwave assisted synthesis and evaluation of antimicrobial activity of 3-{3-(*s*-aryl and *s*-heteroaromatic) acryloyl}-2*H*-chromen-2-one derivatives. *J Heterocycl Chem* 2010;47:179–87.
- [27] Ajani OO, Obafemi CA, Nwinyi OC, Akinpelu DA. Microwave assisted synthesis and antimicrobial activity of 2-quinoxalinone-3-hydrazone derivatives. *Bioorg Med Chem* 2010;18(1):214–21.
- [28] Ajani OO, Aderohunmu DV, Olorunshola SJ, Ikpo CO, Olanrewaju IO. Facile synthesis, characterization and antimicrobial activity of 2-alkanamino benzimidazole derivatives. *Orient J Chem* 2016;32(1):109–20.
- [29] Rithe SR, Jagtap RS, Ubarhande SS. One pot synthesis of substituted benzimidazole derivatives and their characterization. *Rasayan J Chem* 2015;8:213–7.
- [30] Komurcu SG, Rollas S, Uglen M, Gorrod JW. Evaluation of some aryl hydrazones of *p*-aminobenzoic acid hydrazide as antimicrobial agents and their *in-vitro* hepatic microsomal metabolism. *Boll Chim Farmac* 1995;134:375–9.
- [31] Prescott LM, Harley JP, Donald KA. In microbiology. 6th ed. McGraw Hill; 2005.
- [32] Nwinyi OC, Chinedu SN, Ajani OO, Ikpo CO, Ogunniran KO. Antibacterial effects of extracts of *Ocimum gratissimum* and *Piper guineense* on *Escherichia coli* and *Staphylococcus aureus*. *Afr J Food Sci* 2009;3(3):77–81.
- [33] Nwinyi OC, Chinedu NS, Ajani OO. Evaluation of antibacterial activity of *Pisidium guajava* and *Gongronema latifolium*. *J Med Plant Res* 2008;2(8):189–92.
- [34] Ajani OO, Familoni OB, Wu F, Echeme JO, Sujiang Z. Room temperature synthesis and antibacterial activity of new sulfonamides containing *N,N*-diethyl-substituted amido moieties. *Int J Med Chem* 2012;2012:13. doi: <https://doi.org/10.1155/2012/36781>. Article ID 367815.